

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	7.4 (6.2-8.9)	4,968	9.1 (6.5-12.5)	630			●
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	16.4 (15.4-17.6)	4,923	20.1 (16.9-23.8)	619		●	
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	5.0 (4.0-6.1)	2,876	8.2 (5.4-12.2)	305			●
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	35.2 (32.8-37.7)	2,924	33.4 (27.7-39.6)	311			●
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	13.2 (11.9-14.7)	4,909	19.0 (15.6-23.0)	622		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	2.5 (2.1-3.0)	4,911	4.9 (3.2-7.4)	624		●	
QN15: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	8.8 (7.0-10.9)	4,952	16.9 (13.6-20.8)	632		●	
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	7.4 (6.5-8.5)	4,954	10.9 (8.7-13.6)	632		●	
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	19.9 (18.4-21.5)	4,912	28.3 (24.4-32.5)	618		●	
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	6.8 (5.9-7.8)	4,912	12.4 (9.8-15.7)	617		●	
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	4.5 (3.8-5.4)	4,894	17.2 (14.1-20.8)	620		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)	7.5 (6.6-8.5)	4,869	22.8 (19.1-27.0)	599		●	
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	4.6 (3.9-5.5)	3,052	14.0 (10.9-17.8)	394		●	
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	6.1 (5.3-7.1)	3,136	20.4 (15.5-26.3)	425		●	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	12.0 (11.0-13.2)	4,963	27.5 (23.2-32.2)	625		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Florida High School Survey
Risk Behaviors and Sexual Identity Report**

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	9.4 (8.4-10.5)	4,961	25.4 (21.6-29.6)	623		●	
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	23.3 (22.0-24.8)	4,943	57.8 (53.0-62.5)	626		●	
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	9.8 (9.0-10.7)	4,946	39.5 (35.5-43.7)	624		●	
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	7.1 (6.4-7.9)	4,927	31.6 (27.6-35.9)	617		●	
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	4.7 (4.1-5.5)	4,317	22.5 (18.5-27.2)	579		●	
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	1.3 (1.0-1.7)	4,229	6.8 (4.8-9.6)	566		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)	16.7 (15.2-18.2)	4,786	33.1 (29.1-37.3)	598		●	
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)	5.7 (5.0-6.6)	4,851	11.6 (9.2-14.5)	612		●	
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	4.6 (3.8-5.5)	4,832	12.7 (10.5-15.4)	601		●	
QNFRCIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	0.9 (0.7-1.2)	4,832	2.9 (1.6-5.1)	601		●	
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.8 (0.6-1.1)	4,832	1.2 (0.6-2.6)	601			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN40: Percentage of students who ever drank alcohol (at least one drink of alcohol, on at least 1 day during their life)	55.3 (53.3-57.2)	4,599	71.5 (67.6-75.2)	571		●	
QN41: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	15.7 (14.4-17.2)	4,616	22.4 (18.4-27.0)	586		●	
QN42: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	26.0 (24.3-27.8)	4,754	37.5 (33.3-41.9)	592		●	
QN44: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	11.6 (10.3-13.0)	4,883	20.6 (17.0-24.8)	608		●	
QN46: Percentage of students who ever used marijuana (one or more times during their life)	33.1 (31.1-35.2)	4,821	49.9 (46.3-53.5)	595		●	
QN47: Percentage of students who tried marijuana for the first time before age 13 years	6.3 (5.4-7.4)	4,891	11.8 (8.9-15.5)	616		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN48: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	19.0 (17.5-20.5)	4,910	30.7 (27.8-33.7)	610		●	
QN49: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	3.3 (2.7-4.0)	4,963	9.7 (7.3-12.9)	626		●	
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)	2.6 (2.0-3.3)	4,892	8.1 (5.8-11.1)	611		●	
QN56: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	8.9 (7.9-10.0)	4,963	23.8 (20.5-27.4)	625		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	15.4 (14.1-16.9)	4,953	27.3 (23.5-31.4)	623		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	37.1 (34.9-39.3)	4,683	48.4 (44.9-51.9)	582		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	4.4 (3.6-5.2)	4,662	7.4 (5.1-10.6)	581		●	
QN61: Percentage of students who had sexual intercourse with four or more persons during their life	9.4 (8.4-10.5)	4,669	12.7 (10.0-16.1)	579		●	
QN62: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	25.8 (24.3-27.4)	4,674	31.5 (28.2-35.0)	580		●	
QN63: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	18.4 (15.9-21.2)	1,146	22.9 (16.5-31.0)	181			●
QN64: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	39.0 (36.0-42.1)	1,115	58.9 (51.3-66.2)	174		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN65: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	83.0 (80.3-85.4)	1,090	86.1 (78.0-91.5)	162			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	98.5 (97.6-99.1)	1,090	98.5 (95.4-99.6)	162			●
QNSHPARG: Percentage of students who did not use a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	96.6 (95.3-97.6)	1,090	95.9 (88.9-98.6)	162			●
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	78.1 (75.3-80.7)	1,090	80.5 (70.7-87.7)	162			●
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	92.1 (90.4-93.5)	1,084	95.8 (88.6-98.5)	161			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	10.7 (9.1-12.6)	1,090	29.1 (21.9-37.7)	162		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]	10.4 (9.1-11.9)	4,005	14.2 (10.7-18.5)	513			●
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]	13.8 (12.7-15.0)	4,005	14.3 (11.7-17.5)	513			●
QN68: Percentage of students who described themselves as slightly or very overweight	27.9 (26.3-29.6)	4,938	41.9 (38.1-45.8)	627		●	
QN69: Percentage of students who were not trying to lose weight	58.2 (56.4-60.1)	4,944	48.2 (43.9-52.6)	627	●		
QN70: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	28.1 (26.6-29.7)	4,918	26.5 (23.1-30.2)	623			●
QN71: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	12.9 (11.8-14.0)	4,916	16.1 (13.2-19.6)	621		●	

*Confidence interval.

[†]Based on t-test analyses, $p < 0.05$.

[§]Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	6.8 (6.0-7.8)	4,908	7.5 (5.9-9.6)	619			●
QN72: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	41.4 (39.4-43.4)	4,912	40.0 (35.9-44.2)	622			●
QN73: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)	39.7 (38.0-41.4)	4,910	40.6 (35.9-45.5)	624			●
QN74: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)	61.3 (59.5-63.1)	4,895	57.9 (54.0-61.7)	620			●
QN75: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)	24.2 (22.6-25.9)	4,886	22.4 (19.0-26.2)	618			●
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	9.9 (9.0-11.0)	4,861	8.2 (6.0-11.1)	617			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	17.1 (15.6-18.7)	4,809	18.9 (15.9-22.4)	603			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	10.7 (9.6-11.9)	4,809	14.5 (11.9-17.6)	603		●	
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	6.2 (5.3-7.2)	4,809	8.5 (6.6-10.9)	603		●	
QN77: Percentage of students who did not drink milk (during the 7 days before the survey)	28.5 (27.1-30.1)	4,785	32.6 (28.6-36.9)	603			●
QN78: Percentage of students who did not eat breakfast (during the 7 days before the survey)	15.6 (14.1-17.2)	4,799	24.9 (20.6-29.6)	604		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	61.4 (58.7-63.9)	4,799	76.8 (72.5-80.7)	604		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN79: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	58.4 (56.5-60.4)	4,858	72.6 (68.5-76.4)	619		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	20.9 (19.3-22.5)	4,858	26.5 (22.5-30.8)	619		●	
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	75.8 (74.3-77.3)	4,858	85.0 (81.4-87.9)	619		●	
QN80: Percentage of students who watched television 3 or more hours per day (on an average school day)	23.0 (21.2-25.0)	4,865	24.6 (20.8-28.9)	617			●
QN81: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	43.3 (41.4-45.3)	4,847	57.2 (52.7-61.6)	617		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN82: Percentage of students who did not attend physical education (PE) classes on 1 or more days (in an average week when they were in school)	58.1 (55.2-61.0)	4,768	63.7 (58.8-68.3)	600		●	
QNDLYPE: Percentage of students who did not attend physical education classes on all 5 days (in an average week when they were in school)	77.2 (74.0-80.1)	4,768	83.6 (78.9-87.4)	600		●	
QN83: Percentage of students who did not play on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)	51.5 (49.3-53.6)	4,845	61.8 (57.4-66.0)	617		●	
QN84: Percentage of students who had a concussion from playing a sport or being physically active (one or more times during the 12 months before the survey)	12.0 (10.7-13.6)	4,791	13.5 (10.7-16.9)	605			●
QN88: Percentage of students who did not get 8 or more hours of sleep (on an average school night)	78.5 (77.0-80.0)	4,821	83.0 (78.8-86.5)	613		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN85: Percentage of students who were never tested for human immunodeficiency virus (HIV) (not including tests done when donating blood)	88.6 (86.9-90.0)	4,848	84.0 (80.5-87.0)	615	●		
QN86: Percentage of students who did not see a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)	31.4 (29.0-34.0)	4,836	38.8 (35.5-42.2)	612		●	
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)	4.1 (3.3-5.1)	4,836	5.0 (3.2-7.6)	612			●
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	21.1 (19.6-22.8)	4,835	28.2 (24.9-31.7)	613		●	
QN89: Percentage of students who described their grades in school as mostly C's, D's, or F's (during the 12 months before the survey)	24.9 (22.8-27.1)	4,757	34.0 (29.8-38.4)	608		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN90: Percentage of students who used a handheld device to talk or use apps while driving a car or other vehicle (such as the internet, FaceTime, G-mail, Twitter, Facebook, and Snapchat, on at least 1 day during the 30 days before the survey, among students who drove a car or other vehicle during the 30 days before the survey)	38.5 (36.0-41.0)	2,880	39.3 (32.9-46.1)	310			●
QN91: Percentage of students who reported that their property had been stolen or deliberately damaged on school property one or more times (such as their car, clothing, or books, during the 12 months before the survey)	17.3 (16.1-18.6)	4,642	24.2 (20.3-28.6)	589		●	
QN92: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	3.0 (2.4-3.6)	4,655	6.4 (4.2-9.7)	589		●	
QN93: Percentage of students who reported that someone they were dating or going out with threatened them, limited their activities against their will, or made them feel unsafe in any other way (one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the	7.4 (6.4-8.6)	3,032	20.5 (16.9-24.6)	406		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN94: Percentage of students who have been the victim of teasing or name calling because of their weight, size, or physical appearance (during the 12 months before the survey)	19.2 (17.9-20.5)	4,750	39.1 (34.7-43.8)	607		●	
QN95: Percentage of students who have been the victim of teasing or name calling because someone thought they were gay, lesbian, or bisexual (during the 12 months before the survey)	5.0 (4.1-5.9)	4,797	41.9 (37.6-46.4)	605		●	
QN96: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	9.6 (8.8-10.4)	4,687	41.6 (37.6-45.7)	588		●	
QN97: Percentage of students who have been pregnant or gotten someone pregnant (one or more times)	2.6 (2.0-3.3)	4,639	5.3 (3.8-7.4)	586		●	
QN98: Percentage of students who had oral sex	35.9 (33.9-37.9)	4,698	51.3 (47.6-55.1)	598		●	
QN101: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	9.4 (8.3-10.6)	4,613	25.7 (22.4-29.2)	585		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN102: Percentage of students who took some diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight (during the 30 days before the survey)	4.6 (3.9-5.5)	4,687	8.2 (6.0-11.1)	591		●	
QN103: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	3.1 (2.6-3.7)	4,619	11.0 (8.8-13.5)	584		●	
QN104: Percentage of students who drank a can, bottle, or glass of a sports drink (such as Gatorade or PowerAde, not counting low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	54.3 (52.3-56.2)	4,633	44.4 (40.1-48.8)	592	●		
QN105: Percentage of students who drank a can, bottle, or glass of an energy drink (such as Red Bull or Jolt, not including diet energy drinks or sports drinks such as Gatorade or PowerAde, one or more times per day during the 7 days before the survey)	4.0 (3.4-4.7)	4,636	6.6 (4.7-9.2)	592		●	
QN106: Percentage of students who ate at least one meal or snack from a fast food restaurant (such as McDonald's, Taco Bell, or KFC, on three or more days during the 7 days before the survey)	28.4 (26.6-30.3)	4,627	31.5 (27.8-35.5)	590			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN107: Percentage of students who ate dinner at home with at least one of their parents on less than four days (during the 7 days before the survey)	33.3 (31.2-35.4)	4,591	42.6 (38.7-46.5)	587		●	
QN108: Percentage of students who reported their doctor or nurse had not discussed ways to prevent pregnancy, AIDS, or other sexually transmitted diseases (STDs) or they were not sure if they had (during their last check-up, among students who have had a check-up)	65.4 (63.6-67.1)	4,033	59.3 (56.3-62.2)	517	●		
QN109: Percentage of students who have not talked or are not sure if they have talked about AIDS or HIV infection with their parents or other adults in their family	58.5 (56.9-60.2)	4,539	56.5 (52.4-60.5)	576			●
QN110: Percentage of students who have not ever been taught or are not sure if they have ever been taught about AIDS or HIV infection during grades 9 through 12	24.7 (22.8-26.7)	4,546	27.2 (23.3-31.5)	571			●
QN111: Percentage of students who think it is somewhat important or not important for schools to help students address the problems of today (such as drug abuse, violence, AIDS/HIV, teen pregnancy, abuse, and suicide)	11.7 (10.6-13.0)	4,538	11.0 (8.0-14.8)	572			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN113: Percentage of students who reported that, if they or one of their friends were being physically or sexually abused, they do not know or are not sure who they should report it to	24.4 (22.9-26.0)	4,531	23.4 (19.8-27.4)	572			●
QN114: Percentage of students who never, rarely or sometimes wear sunscreen (with an SPF of 15 or higher when they are outside for more than one hour on a sunny day)	91.5 (90.4-92.5)	4,516	90.2 (86.9-92.8)	574			●
QN115: Percentage of students who have not received a flu shot (during the 12 months before the survey)	44.3 (42.5-46.1)	4,521	50.8 (45.9-55.6)	574		●	
QNSPDRK1: Percentage of students who drank a can, bottle, or glass of a sports drink one or more times per day (such as Gatorade or Powerade, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	14.4 (13.0-15.9)	4,633	11.6 (8.8-15.2)	592			●
QNSPDRK2: Percentage of students who drank a can, bottle, or glass of a sports drink two or more times per day (such as Gatorade or Powerade, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	8.6 (7.6-9.8)	4,633	6.7 (4.8-9.4)	592			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Florida High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSPDRK3: Percentage of students who drank a can, bottle, or glass of a sports drink three or more times per day (such as Gatorade or Powerade, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	4.9 (4.1-5.8)	4,633	3.2 (1.9-5.5)	592			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.